
Comparison of the Effect of Miconazole and Clotrimazole in the Treatment of Vulvovaginal Candidiasis Among Women Seen in a Tertiary Medical Center from 2016 to 2020*

Shiara Marriz T. Marquez, MD and Lylah D. Reyes, MD, FPOGS

ABSTRACT

Vulvovaginal candidiasis (VVC) is one of the frequent infections of the female genital tract and is the second most common cause of vaginal infections after bacterial vaginosis. According to the Center for Disease Control and Prevention (CDC), azoles are the first-line of treatment for VVC. Among the azoles available in the Philippines, only miconazole and clotrimazole are recommended for both pregnant and non-pregnant women. This study reviewed a total of 316 records of patients, regardless of pregnancy status who were given either single dose 1200 mg miconazole or a 6-day 100 mg clotrimazole vaginal suppository as treatment for vulvovaginal candidiasis. Upon follow-up, 11 out of 316 (3.46%) were still symptomatic which was noted in 2 (18.1%) from the miconazole group and 9 (81.8%) from the clotrimazole group. The difference for these outcomes between the treatment groups is statistically significant. The treatment failure rate for miconazole is 1.2% (2/161) and 5.8% (9/155) for clotrimazole. None of the charts of these women had recorded adverse reaction to the given treatment. In conclusion, the single-dose miconazole regimen has a significantly higher clinical cure rate than the 6-day clotrimazole. It has the potential to improve patient compliance and improve treatment outcome at a lower cost.

Key words: vulvovaginal candidiasis, miconazole, clotrimazole, azole antifungals, vaginitis

Vaginitis is one of the most common reasons why women seek consult from obstetrician and gynecologists. Vulvovaginal candidiasis is one of the frequent infections of the female genital tract and is the second most common cause of vaginal infections after bacterial vaginosis. More than 90% of cases are caused by *Candida albicans*.¹ It affects 75% of women at least once in their lifetime, with 50% developing a second episode. About 5% of women will have recurrent vulvovaginal candidiasis, defined as four or more episodes of such infection within a year.²

Vulvovaginal candidiasis (VVC) is usually diagnosed based on clinical signs and symptoms or microscopic examination, or a combination of both. With regards to treatment of VVC, based on

Center for Disease Control and Prevention (CDC) recommendation, azoles are considered as the first-line drug class. It is further categorized as uncomplicated VVC which involves a short course of azole antifungals and complicated VVC wherein azoles are given for a longer duration. However, due to poor compliance, increasing recurrence rates and emergence of drug-resistant *Candida* strains worldwide, further studies are being done to determine which among the recommended drugs is the ideal regimen.

The azoles that are available in the Philippines include clotrimazole, miconazole and fluconazole. Miconazole and clotrimazole are recommended for both pregnant and non-pregnant women. However, only Miconazole can be given as a single dose. Hence, in terms of management, single dosing regimen may translate to better compliance than multiple dosing regimen. This study, therefore, would like to compare the effect of the 6-day 100

* From the Department of Obstetrics and Gynecology

mg Clotrimazole suppository versus the single-dose 1200 mg Miconazole suppository in the treatment of vulvovaginal candidiasis.

Review of Related Literature

The CDC classifies vulvovaginal candidiasis into two: uncomplicated and complicated vulvovaginal candidiasis.³ Uncomplicated vulvovaginal candidiasis includes infections caused by *Candida albicans*, periodic vulvovaginal candidiasis, mild to moderate vulvovaginal candidiasis and those affecting a non-immunocompromised host. On the other hand, complicated vulvovaginal candidiasis encompasses infections caused by non-*albicans* *Candida*, recurrent vulvovaginal candidiasis, or severe vulvovaginal candidiasis, or those affecting immunocompromised hosts.

Signs and symptoms include the presence of a thick, white vaginal discharge in association with vulvar itching, burning with or without dysuria. Most women with signs and symptoms of vulvovaginal candidiasis can be diagnosed with the use of saline wet mount, 10% potassium hydroxide (KOH) test, or gram stain of vaginal discharge. In cases with VVC, the smear demonstrates hyphae or pseudohyphae of *Candida* species, or a vaginal culture test shows a *Candida* species.⁴

Age appears to be an important risk factor in the overall incidence of vulvovaginal candidiasis because episodes occur mostly during the reproductive years.⁴ Other factors like pregnancy, diabetes mellitus, immunosuppression, use of artificial contraceptives (pills, intrauterine devices, diaphragms), poor perineal hygiene, and vaginal douching are also contributory for the development of vulvovaginal candidiasis. Aside from the fact that poor compliance to treatment may also be a contributing factor for recurrence. Hence, if treatment is given as single course with associated improvement in symptomatology and eradication of the organism, issues on compliance with the institution of appropriate treatment are addressed accordingly.

Azole antifungals like miconazole and clotrimazole work by inhibiting the cytochrome P450-dependent enzyme lanosterol 14- α -demethylase that is necessary for the conversion of lanosterol to ergosterol which is a vital component of the cellular membrane of *Candida* species. Their effectiveness in treating vulvovaginal candidiasis has long been a topic in researches.

A computer-based literature search was done to find relevant articles for the study. The search terms vulvovaginal candidiasis, miconazole, clotrimazole; according to rank were used. Union and intersection of the search terms were done. Combination of search terms using vulvovaginal candidiasis, miconazole, clotrimazole yielded 71 articles; among the 71 articles, 10 were relevant to the study. Addition of pregnancy to the combination of search terms yielded 12 articles, 6 of which are relevant to the present study. Hand search was also done thru the Philippine Obstetrical and Gynecological Society database which yielded no article.

The single dose miconazole vaginal suppository consists of 1200 mg of miconazole nitrate along with other inactive ingredients. According to Barnhart, it is proven to exhibit fungistatic and fungicidal activity against *Candida* species. In the same year, the author also conducted a multicenter randomized controlled trial involving 573 women with VVC. Patients were given a single dose 1200 mg miconazole vaginal ovules either in daytime or bedtime.⁵ Clinical cure rates were 74.5% and 73.6% in the daytime and bedtime groups, respectively. The study concluded that the efficacies of daytime and bedtime administration of single dose miconazole were the same.

In a prospective randomized case control study conducted by Fan, et al, a total of 577 cases of VVC were treated with either 1200 mg miconazole vaginal suppository or 150 mg fluconazole tablet and were followed up after treatment.⁶ After seven days of treatment with miconazole, the clinical cure rate is 75.9% while the mycological cure rate is 84%. The study concluded that therapy with intravaginal miconazole is as effective as fluconazole in treating vulvovaginal candidiasis.

Two randomized, single-blind, multicenter, controlled trials involving 558 patients with VVC were conducted by Upmalis, et al. in the year 2000.⁷ They tried to determine the efficacy of single dose 1200 mg miconazole vaginal ovule versus seven consecutive doses of miconazole 2% vaginal cream. In both studies, patients were assigned to two treatment groups and were followed up 14 to 18 days after the initiation of treatment. For the first study, clinical cure rate was 93% for the miconazole ovule and 97% for the miconazole cream. While for the second study, clinical cure rate was 88% for the miconazole ovule and 84% for the miconazole cream. They concluded that single dose miconazole ovule is as efficacious

as 7-day miconazole cream in the treatment of vulvovaginal candidiasis.

Clotrimazole is a broad-spectrum antimycotic drug that is also widely used in treating vulvovaginal candidiasis. The intravaginal forms are either 100, 200 or 500 mg, given daily for 6, 3 or 1 day(s) respectively.⁸

In an open, parallel randomized study conducted by Zhou, et al. there were 240 women with VVC who were either given 2 doses of 500 mg Clotrimazole intravaginally or 2 doses of 150 mg Fluconazole orally.⁹ After 7 to 14 days, the clinical cure rate in the clotrimazole group was 88.7% and 89.1% in the fluconazole group. The researchers concluded that 2 doses of 500 mg Clotrimazole given intravaginally is as effective as 2 doses of 150 mg oral fluconazole in treating VVC.

Sekhavat, et al. conducted a prospective study by involving 142 patients with acute vulvovaginal candidiasis.¹⁰ Participants were either given oral single-dose fluconazole tablet or 200 mg clotrimazole intravaginal tablet for 6 days. Patients were followed up 7 days and 1 month post treatment. After 7 days, 61 patients (84.7%) were cured clinically and 58 patients (80.5%) mycologically in the fluconazole group and 60 patients (83.3%) were cured clinically and 49 patients (70%) mycologically in the clotrimazole group. The authors concluded that both treatment regimen have comparable clinical efficacy.

Attempts to compare the efficacy of miconazole and clotrimazole can be traced back in the 1970s. Eliot, et al. conducted a multicenter randomized controlled trial comparing the efficacy of nystatin, clotrimazole and miconazole in the treatment of 166 women with VVC.¹¹ Patients were given either 100, 000 units nystatin pessaries nightly for 14 days, 100 mg clotrimazole vaginal tablets nightly for 6 nights, 100 mg miconazole pessaries nightly for 14 days or 100 mg miconazole pessaries in the morning and in the evening for 7 days. Mycological cure rate 4 weeks post treatment was 93% for nystatin, 88% for clotrimazole, 91% for 14-day miconazole and 86% for the 7-day miconazole group. The trial concluded that cure rates were comparable for the 4 treatment groups. However, relapse rates were higher for those in the clotrimazole and nystatin groups at 50% and 28.3% respectively compared with the 14-day miconazole (2.4%) and 7-day miconazole groups (3.4%).

Balsdon (1981) conducted a randomized controlled trial comparing the use of twice daily 100 mg miconazole-coated tampons for 5 days and six-

day course of clotrimazole intravaginal tablets once a day in the treatment of VVC in 100 women.¹² Patients were followed up 7 days after treatment where clinical cure was found to be 98% for clotrimazole and 99% for the miconazole group. The study concluded that both treatment regimens were equally effective in treating VVC. However, the miconazole coated tampons were more acceptable to patients because of the lesser duration of treatment and ease of use, as there was no leakage of medications from the vagina.

Common adverse reactions with the use of azole antifungals include vomiting, loose stools, abdominal pain, paresthesia, headache, dizziness, fever, chills, vaginal burning, stinging, itching and irritation.³ More of these systemic adverse effects are likely to be reported with oral compared with intravaginal antifungal administration.⁴ In the same study by Fan, et al., adverse effect rate of miconazole was found to be at 0.9%.⁶ The adverse effects reported were vulvar burning, pain, edema and itching. Moreover, reports on the adverse effects of clotrimazole were mainly local and similar to that of miconazole. In the previously mentioned study of Zhou, et al., 16 out of 115 patients who received vaginal clotrimazole experienced adverse events, most of which include vulvovaginal pruritus, burning, irritation and bleeding.⁹

Topical antifungals like miconazole and clotrimazole are classified as Pregnancy Category C. They have minimal absorption hence can be given at anytime during pregnancy.¹³ A retrospective cohort study by Rotem et al. involving 2,306 pregnancies evaluated the risk for major malformations following first- trimester exposure to vaginal azoles (clotrimazole and miconazole).¹⁴ Crude and adjusted relative risks for major congenital malformations and for specific malformations according to organ systems were calculated using a multivariate negative binomial regression. They found no significant association between first trimester exposure to clotrimazole nor miconazole vaginal tablets and total major malformations or specific malformations according to organ systems.

Daniel, et al. also conducted a historical cohort study involving 3246 pregnancies with exposure to vaginal antimycotics before 20 weeks age of gestation for treatment of vulvovaginal candidiasis.¹⁵ Clotrimazole was administered to 2,712 women while 633 women received miconazole. A computerized database of medication dispensation was linked with

2 computerized databases containing information on births and spontaneous abortions. Time-varying Cox regression models were constructed adjusting for mother's age, diabetes mellitus, hypothyroidism, obesity, hypercoagulable or inflammatory conditions, recurrent miscarriages, intrauterine contraceptive device, ethnicity, tobacco use, and the year of admission. They concluded that exposure to vaginal antimycotics such as miconazole and clotrimazole was not associated with spontaneous abortion (Adjusted hazard ratio=1.34; 95% CI, 0.99-1.8).

Based on the available evidence both miconazole and clotrimazole are efficacious in the treatment of vulvovaginal candidiasis. However, in terms of studies comparing the efficacy of these two antifungals, they were found to be outdated (1979-1981). Even so, there are still no studies based on computerized literature search comparing the effect of the single dose miconazole regimen with azoles that can be given regardless of pregnancy status like clotrimazole. A meta-analysis conducted by Qin, et al. reviewed 41 randomized controlled trials on the efficacy of the different antifungal drugs used in the treatment of vulvovaginal candidiasis.¹⁶ None of them compared the use of single dose miconazole with clotrimazole. Hence, in the present study, the authors would like to assess the effect of the single dose 1200 mg miconazole compared with 6-day 100 mg clotrimazole regimen in treating vulvovaginal candidiasis.

Significance of Study

More and more women, especially those who are pregnant, are troubled by the high prevalence and recurrence rates of vulvovaginal candidiasis. In this institution, the most commonly prescribed azole is clotrimazole followed by miconazole. Although azoles are efficacious in providing complete cure for vulvovaginal candidiasis, only miconazole can be given as a single-dose for the treatment of this infection. The findings in this study will primarily have implications for ensuring compliance of patients. The advantage of single dose regimen is that treatment is more convenient hence, subsequently improving compliance. Thereby, the primary implication of the findings in this study is toward ensuring compliance of patients. This can be contributory in promoting appropriate treatment that will consequently lower the incidence and associated healthcare cost in the management of vulvovaginal candidiasis. Ultimately,

this will translate to improved health outcomes among women with vulvovaginal candidiasis regardless of pregnancy status.

The objective of the study was to compare the effect of miconazole versus clotrimazole in the treatment of vulvovaginal candidiasis among patients seen at the out-patient department.

MATERIALS AND METHODS

Study Design

This is a retrospective cohort study.

Setting and Target Population

This study entailed review of records of patients at the Obstetrics and Gynecology Out-patient department of this tertiary private medical center from January 2016 to March 2020. This study included women regardless of pregnancy status who were given either intravaginal miconazole or clotrimazole as treatment for vulvovaginal candidiasis.

Inclusion Criteria

All women fulfilling the following criteria were included in this study:

1. Age 18 years to 35 years old
2. Those with ANY ONE of the following signs and symptoms of vulvovaginal candidiasis:
 - 2.1.Curd-like vaginal discharge
 - 2.2.Vulvar pruritus
 - 2.3.Dyspareunia
 - 2.4 Vulvar erythema
3. Those with pseudohyphal, or hyphal elements seen on Gram Stain, Wet Smear or KOH
4. Those who completed the treatment and had follow-up evaluation recorded in the chart

Exclusion Criteria

Those women with the following conditions were excluded:

1. Current infection or infection within 2 weeks from recruitment involving other types or

causes of vaginitis such as bacterial vaginosis and trichomoniasis as well as those with pelvic inflammatory disease and pyelonephritis

2. Complicated vulvovaginal candidiasis (four or more episodes of candidiasis per year (recurrent vulvovaginal candidiasis or severe symptoms or findings (severe VVC), or non-albicans *Candida* infection, or abnormal host (e.g. uncontrolled diabetes, debilitation or immunosuppression).

Methodology

This study involved review of charts for a period of 4 years from 2016 to 2020. The records were gathered from the medical records section of this tertiary hospital after a consent from the chief information and data privacy officer was secured. The eligible records underwent data extraction by the primary investigator. Gathered data were encoded in a standard data collection sheet. The data included the patient's age, gravidity and parity, signs and symptoms (curd-like vaginal discharge, vulvar pruritus, dyspareunia, vulvar erythema) on initial consult and its persistence or resolution on follow-up, presence of pseudohyphal, hyphal, or fungal elements seen on Gram Stain, Wet Smear or KOH, the antifungal agent used, route of administration, its associated complications as well as adverse effects. Purposive sampling was done for this study.

Sample Size Calculation

The sample size was computed based on the formula of comparison of proportions from two independent samples. These proportions were based on two studies: 1) Fan, et al. reported that 75.9% of their participants given single dose Miconazole for vulvovaginal candidiasis was noted to have clinical cure⁶, 2) Zhou, et al. found that the clinical cure rate among women with VVC given clotrimazole is 88.7%.⁹

With a proportion, P1, of 0.75 in group 1 and P2 of 0.88 in group 2, a total of 308 charts of women seen at the outpatient department, about 154 for each treatment group was reviewed in the study. This sample size gives a probability of 80% of rejecting the null hypothesis of equal proportions if the alternative holds. The test of equality of proportions was carried out at 0.05 level of significance.

Data Management and Analysis

Microsoft excel was used for data entry and encoding. For data analysis, Stata version 9.0 was used. Univariate analysis such as mean and range was used to describe continuous variables such as age, gravidity and parity. Frequency distribution was used to describe the categorical variables which include the proportion of participants with presence or absence of symptoms as well as the pseudohyphal, or hyphal elements on microbiologic examination. The presence or absence of side effects was also described using frequency distribution. Chi square was used to analyze the comparison of the proportion of the participants with the above categorical variables, while t-test was used to compare the continuous variables between the two treatment groups.

Ethical Consideration

The study proposal before its implementation underwent review and was approved by the technical and ethics review board of the Research Center for Development. Permission to retrieve and review the records was obtained from the office of the chief medical officer, data privacy officer, and the medical records section.

This study complied with the Data Privacy Act of 2012. Pertinent information such as name and address were not included nor mentioned in this study. Each patient was assigned a specific number code to represent her case on the data collection tool. No identifiable information pertaining to the participant appeared on the data collection sheet. These ensured anonymity of the participants in this study as well as maintained confidentiality.

A waiver of informed consent was obtained from the board of ethics. There was no direct risk involved since this only entailed review of chart records of the eligible participants. The personal data taken from patient charts were not included so that no identifiable information pertaining to the patient will appear in any part of the paper. There were no conflicts of interest involved in this study. The authors have no affiliations with or involvement in any organization or entity with any financial or non-financial interest in the subject matter or material discussed in this manuscript.

RESULTS

A total of 316 charts were eligible to be included in this study. Among women whose chart records

were included, there were 161 (50.94%) who used miconazole while 155 (49.05%) took clotrimazole. The age range of women whose charts were included were 18 to 35 years old with a mean age of 28. Among the chart records, 16% (53/316) were pregnant and 82% (263/316) were non-pregnant. The range of gravidity was from 0 to 5 with a mean gravidity of 1 while the range of parity was 0 to 3 with a mean parity of 1. However, when the baseline characteristics of women were compared, the record showed that those women in the miconazole group were significantly older with greater parity as compared to those given clotrimazole. (Table 1)

All of the 316 records of women included in the study presented with signs and symptoms of vulvovaginal candidiasis. The most common of which is curd-like vaginal discharge (86.7%), followed by vulvar pruritus (80.7%), vulvar erythema (50.6%), and the least occurring is dyspareunia (0.94%). When the signs and symptoms were individually compared between the two treatment arms, no statistically significant differences were noted. Table 2 shows the detailed comparison on the different signs and symptoms between the two treatment arms.

During the chart review, women diagnosed with vulvovaginal candidiasis were either given single dose 1200 mg miconazole or 6-day 100 mg clotrimazole and were asked to follow up after 7 days from initiation of treatment. Upon follow-up, 11 out of 316 (3.46%) were still symptomatic which was noted in 2 (18.1%) from the miconazole group and 9 (81.8%) from the clotrimazole group. (See table 3 for details). The difference for these outcomes between the treatment groups is statistically significant. The treatment failure rate is 1.2% (2/161) for miconazole and 5.8% (9/155) for clotrimazole.

In this study, adverse effects of intravaginal antifungal treatment such as vulvar burning, pain, edema and itching were also reviewed. However, none of the charts of these women had recorded adverse reaction to the given treatment.

DISCUSSION

Vulvovaginal candidiasis remains one of the most common reasons for women to seek consult to the clinics. According to Al Quaiz, around 3% of women

Table 1. Comparison of baseline characteristics as to age, gravidity, parity and pregnancy status between miconazole and clotrimazole

Baseline Characteristics	Antifungal Agent		p value
	Miconazole Mean ($\pm$ SD)	Clotrimazole Mean ($\pm$ SD)	
Age	29.78 (3.49)	28.04 (4.04)	0.001
Gravidity	1.15 (0.96)	0.95 (0.84)	0.05
Parity	0.84 (0.82)	0.63 (0.95)	0.01
Pregnancy status n(%)			0.33
Pregnant	23 (14)	30 (19)	
Non-pregnant	138 (55)	125 (80)	

Table 2. Comparison of baseline signs and symptoms between miconazole and clotrimazole.

Baseline signs and symptoms	Antifungal Agent		p value
	Miconazole n (%)	Clotrimazole n (%)	
Curd-like vaginal discharge	135 (83)	139 (89)	0.13
Pruritus	127 (78)	128 (82)	0.40
Dyspareunia	0	3 (1)	0.08
Erythema	82 (50)	78 (50)	0.91

Table 3. Comparison of the resolution of signs and symptoms with treatment between miconazole and clotrimazole

Signs and Symptoms of vulvovaginal candidiasis	Antifungal Agent		p value
	Miconazole N=161 n (%)	Clotrimazole N=155 n (%)	
Absent	159 (52.4)	146 (47.5)	0.027
Present	2 (18.1)	9 (81.8)	

presenting with signs and symptoms of vulvovaginal candidiasis usually consult at the clinic.¹⁷

Among all the antifungal agents used in treating vulvovaginal candidiasis, only miconazole and clotrimazole are proven safe for both pregnant and non-pregnant women. The intravaginal forms of these antifungals allow effective treatment of the disease with the least side effects possible. Clotrimazole is given as a 6-day regimen but only Miconazole can be given as a single dose treatment.

The present study compared the effect of miconazole versus clotrimazole in the treatment of vulvovaginal candidiasis among patients seen at the out-patient department of this tertiary hospital from 2016 to 2020. The result of the comparison between the two types of antifungal agents revealed that single dose 1200 mg miconazole is more efficacious than the 6-day 100 mg clotrimazole regimen in the promoting clinical cure of vulvovaginal candidiasis. The clinical cure rate for miconazole is 98% and 94% for clotrimazole. Thereby, treatment failure rate as expected was lower among those in the miconazole group (1.2%) as compared to those in the clotrimazole group (5.8%).

In contrast to this study, there were two trials that showed comparable cure rates between miconazole and clotrimazole in the treatment of vulvovaginal candidiasis. The first study was that of Eliot, et al. in 1971, wherein cure rate of the 6-day 100 mg clotrimazole is 88%, 86% for 7-day 100 mg miconazole given twice daily and 91% for 14-day 100 mg miconazole.¹¹ For the second study, which was a randomized controlled trial conducted by Balsdon in 1981, twice daily 100 mg miconazole tampons given for 5 days had a clinical cure rate of 99% while a six-day course of 100 mg clotrimazole intravaginal tablets yielded a clinical cure rate of 98%.¹² Azole antifungals like miconazole and clotrimazole inhibit the cytochrome P450- dependent

enzyme lanosterol 14-alpha-demethylase that is necessary for the synthesis of ergosterol, a vital component of the cellular membrane of *Candida* species.⁹ Several literatures have found out that such enzyme inhibition is dependent on the dose of azoles, with higher concentrations achieving greater fungistatic activity.^{18,19} This is most likely the reason why in this study, the single dose 1200 mg miconazole regimen yielded a higher clinical cure rate than the 6-day 100 mg clotrimazole regimen. Another reason may be due to lesser product misuse since the single dose miconazole regimen only entails one-time application. Lastly, the shorter treatment duration with the single dose miconazole regimen may have contributed to patient compliance during the treatment.

The adverse effects associated with azoles administered intravaginally include vulvar burning, pain, edema and itching.⁴ Such adverse effects are reported by patients given with either miconazole or clotrimazole vaginal suppositories as treatment for VVC.^{6,9} However, according to literatures, the adverse effects of these agents are well- tolerated.⁹ This is probably the reason why in this study, there were no recorded or reported adverse effects during the time of azole treatment. In the study of Fan, et al. there were reports of adverse effects but they were only noted in 2 out of 290 patients who were given intravaginal miconazole.⁶ Similarly, in the study of Zhou, et al. only 16 out of 115 patients developed adverse reactions after administration of intravaginal clotrimazole.⁹ None of them discontinued the treatment because of the said adverse effects. This shows that side effects associated with local antifungal agents remain negligible.

When it comes to cost, the average price of miconazole 1200 mg intravaginal tablet given as a single dose is Php 552.67. In comparison with clotrimazole 100 mg intravaginal tablet, the average

cost is Php 106.50 per suppository usually inserted once a day for 6 days. With this, the total cost for clotrimazole is Php 639.00. Therefore, aside from a shorter treatment duration, the single-dose miconazole is better in terms of affordability with cost savings of Php 86.33 per treatment course. This may translate to better patient compliance that may eventually improve the outcome of treatment.

The main limitation of the present study is its retrospective nature. Cases gathered per arm were not equal. The eligible charts did not include complicated and recurrent vulvovaginal candidiasis since the recorded follow-up were mostly done once, and just immediately after treatment. It may also be useful if such information from patients having these conditions were included. Also, microbiologic cure was not assessed in this study since most of the eligible cases did not have a repeat gram stain, wet smear nor KOH smear of their vaginal discharge post treatment.

CONCLUSION

In this institution, clotrimazole and miconazole are both widely used in the management of vulvovaginal candidiasis. As seen in this study, the single-dose miconazole regimen has a significantly higher clinical cure rate than the 6-day clotrimazole, with adverse effects being uncommon for both agents. Since these agents appear to be comparable in providing relief of symptoms with negligible adverse effects, the single dose miconazole regimen has the potential to improve patient compliance and improve treatment outcome at a lower cost.

RECOMMENDATION

For a more accurate comparison between miconazole and clotrimazole, a prospective randomized controlled study with equal participants per arm is therefore recommended. Future studies may include patients with complicated vulvovaginal candidiasis such as those with comorbidities. Lastly, a more profound investigation may be achieved by assessing the complete cure rate which includes both microbiologic and clinical cure rates between the two treatment arms. It is also worthwhile to determine the recurrence rate of vulvovaginal candidiasis between the two treatment arms. Hence, it is recommended to conduct the trial with a longer follow-up.

REFERENCES

1. Lobo R, Gerhenson D, Lentz G and Valea F. Comprehensive Gynecology 7th edition. Elsevier, Philadelphia, USA. 2017.
2. Martinez R, Franceschini S, Patta M, et al. Improved treatment of vulvovaginal candidiasis with fluconazole plus probiotic *Lactobacillus rhamnosus* GR-1 and *Lactobacillus reuteri* RC-14. The Society for Applied Microbiology, Letters in Applied Microbiology 2009; 48: 269–74.
3. Center for Disease Control and Prevention Sexually Transmitted Diseases Treatment Guidelines (2015). <https://www.cdc.gov/std/tg2015/candidiasis.htm>
4. Xie H, Feng D, Mei L, Chen H, Wang X and Fang F. Probiotics for vulvovaginal candidiasis in non-pregnant women (Review). Cochrane Database of Systematic Reviews 2017; Issue 11. Art. No.: CD010496.
5. Barnhart K. Safety and efficacy of bedtime versus daytime administration of the miconazole nitrate 1200 mg vaginal ovule insert to treat vulvovaginal candidiasis. Curr Med Res Opin. 2005 Jan;21(1):127-34. doi: 10.1185/030079904x18018.
6. Fan S, Liu X, Liang Y. Miconazole nitrate vaginal suppository 1,200 mg versus oral fluconazole 150 mg in treating severe vulvovaginal candidiasis. Gynecol Obstet Invest 2015; 80(2): 113-8. doi: 10.1159/000371759. Epub 2015 Feb 17.
7. Upmalis DH, Cone FL, Lamia CA, Reisman H, Rodriguez-Gomez G, Gilderman L, Bradley L. Single-dose miconazole nitrate vaginal ovule in the treatment of vulvovaginal candidiasis: two single-blind, controlled studies versus miconazole nitrate 100 mg cream for 7 days. J Womens Health Gend Based Med 2000; 9(4): 421-9. doi: 10.1089/15246090050020745.
8. Crowley PD, Gallagher HC. Clotrimazole as a pharmaceutical: past, present and future. J Appl Microbiol 2014; 117(3): 611-7. doi: 10.1111/jam.12554. Epub 2014 Jun 30.
9. Zhou X, Li T, Fan S, Zhu Y, Liu X, Guo X and Liang Y. The efficacy and safety of clotrimazole vaginal tablet vs. oral fluconazole in treating severe vulvovaginal candidiasis. Mycoses 2016; 59: 419- 28. <https://doi.org/10.1111/myc.12485>
10. Sekhavat L, Tabatabaai A, Tezerjani. Oral fluconazole 150 mg single dose versus intravaginal clotrimazole treatment of acute vulvovaginal candidiasis. J Infect Public Health 2011; 4: 195-9.
11. Eliot BW, Howat RC, Mack AE. A comparison between the effects of nystatin, clotrimazole and miconazole on vaginal candidiasis. Br J Obstet Gynaecol 1979; 86(7): 572-7. doi: 10.1111/j.1471-0528.1979.tb10812.x.
12. Baldson MJ. Comparison of miconazole-coated tampons with clotrimazole vaginal tablets in the treatment of vaginal candidosis. Br J Vener Dis 1981; 57(4): 275-8. doi:10.1136/sti.57.4.275
13. Pilms B, Jullien V, Sobel J, Lecuit M, Lortholary O, Charlier C. Antifungal drugs during pregnancy: an updated review. J Antimicrob Chemother 2015; 70(1): 14-22. doi: 10.1093/jac/dku355. Epub 2014 Sep 8.
14. Rotem R, Fishman B, Daniel S, et al. Risk of major congenital malformations following first trimester exposure to vaginal azoles used for treating vulvovaginal candidiasis: a population based retrospective cohort study. BJOG 2018; 125(12): 1550.

15. Daniel S, Rotem R, Koren G, et al. Vaginal antimycotics and the risk for spontaneous abortions. *Am J Obstet Gynecol* 2018; 218: 601.e1-7
16. Qin F, Wang Q, Zhang C, Fang C, Zhang L, Chen H, Zhang M, Cheng F. Efficacy of antifungal drugs in the treatment of vulvovaginal candidiasis: a Bayesian network meta-analysis. *Infect Drug Resist* 2018; 11: 1893-901. doi: 10.2147/IDR.S175588.
17. Al Quaiz J. Patients with vaginal discharge: A survey in a university primary care clinic in Riyadh City. *Ann Saudi Med* 2000. <https://doi.org/10.5144/0256-4947.2000.302>
18. Kjærstad MB, Nielsen F, Nøhr-Jensen L. et al. Systemic uptake of associated enzyme activities in women. *Eur J Clin Pharmacol* 2010; 66: 1189–97.
19. Hitchcock CA, Dickinson K, Brown SB, et al. Interaction of azole antifungal antibiotics with cytochrome P450 dependent 14- α sterol demethylase purified from *Candida albicans*. *Biochem J* 1990; 475-80. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1131156/pdf/biochemj00188-0162.pdf>